

proached them to within 2 m—and they continued feeding on the fruits when we left the site after 15 min of observation.

The fruits of *Flueggia virosa* is eaten by children in Pemba and Zanzibar. In Pemba the shrub is called *mususi mizi*, while on Zanzibar it is known as *mkwamba*. We tasted the ripe berries and found them pulpy and sweet.

Chapin (1954) records instances of berries being found in the stomachs of four species of *Anthreptes* sunbirds and in three out of six stomachs of Olive Sunbirds *N. olivacea* in Zaïre. In southern Africa Maclean (1985) records the Collared Sunbird *Anthreptes collaris* eating small berries.

References

- CHAPIN, J.P. 1954. The birds of the Belgian Congo. Part 4. *Bulletin of the American Museum of Natural History* 75B: 1–846.
- MACLEAN, G.L. 1985. *Roberts' birds of southern Africa*. Cape Town: Trustees of the John Voelcker Bird Book Fund.

A. L. Archer, Box 15676, Nairobi and I. S. C. Parker, Box 15093, Nairobi

Scopus 17: 60–61, June 1993

Received 29 March 1993

A breeding record for the African Spoonbill *Platalea alba* in Tanzania

On 10 August 1990 at Lake Eyasi (35°20'E, 3°26'S) in northern Tanzania, I found a small colony of African Spoonbills *Platalea alba*. The water level in the lake was high for the time of year but was receding quite fast and the small island on which the birds were nesting was only some 100 m from the shore with the surrounding water 60 cm deep.

There were 26 active nests, 14 with young, some of which could walk; the remaining nests contained eggs. Some clutches were still being incubated but a few were probably abandoned. An abandoned egg measured 71.4 x 45.1 mm. A more rounded, less pointed egg found at the nesting site measured 91.6 x 58.6 mm and was probably of a Pink-backed Pelican *Pelecanus rufescens* from an earlier breeding attempt.

Britton (1980) does not list Lake Eyasi as a breeding locality for the African Spoonbill and Brown & Britton (1980) give laying dates as February to April for 250 records and May for one record within their Region C, which includes Lake Eyasi. However, within their Region D, less than 100 km to the west, breeding is "much more elastic" and includes July and August.

References

- BROWN, L.H. & BRITTON, P.L. 1980. *The breeding seasons of East African birds*. Nairobi: EANH.S.
- BRITTON, P.L. 1980 (ED). *Birds of East Africa*. Nairobi: EANH.S.

N. E. Baker, Box 23404, Dar es Salaam, Tanzania

Scopus 17: 61, June 1993

Received 11 October 1991